

Seifert manifolds and $(1, 1)$ -knots

Luigi Grasselli

Michele Mulazzani

Abstract

The aim of this paper is to investigate the relations between Seifert manifolds and $(1, 1)$ -knots. In particular, we prove that every orientable Seifert manifold with invariants

$$\{Oo, 0 \mid -1; \underbrace{(p, q), \dots, (p, q)}_{n \text{ times}}, (l, l-1)\}$$

has the fundamental group cyclically presented by $G_n((x_1^q \cdots x_n^q)^l x_n^{-p})$ and, moreover, it is the n -fold strongly-cyclic covering of the lens space $L(|nlq-p|, q)$, branched over the $(1, 1)$ -knot $K(q, q(nl-2), p-2q, p-q)$, if $p \geq 2q$, and over the $(1, 1)$ -knot $K(p-q, 2q-p, q(nl-2), p-q)$, if $p < 2q$.

2000 Mathematics Subject Classification: Primary 57M12, 57M25; Secondary 20F05, 57M05.

Keywords: Seifert manifolds, $(1, 1)$ -knots, cyclic branched coverings, cyclically presented groups, Heegaard diagrams.

1 Introduction

Cyclic branched coverings of knots in $\mathbf{S}^3$ with cyclically presented fundamental group have been deeply investigated in the recent years by many authors (see [1, 10, 11, 12, 17, 20, 21, 22, 23, 33]). Their results have been included in an organic and more general context in [24], where it is proved that the fundamental group of every n -fold strongly-cyclic branched covering of a $(1, 1)$ -knot admits a cyclic presentation encoded by a genus n Heegaard diagram. In [5]

this result has been improved, obtaining a constructive algorithm which explicitly gives the cyclic presentation, starting from a representation of the $(1, 1)$ -knot via the mapping class group of the twice punctured torus (see [6] for further details on this representation).

In [14], M. J. Dunwoody introduces a class of manifolds depending on six integers, the so-called Dunwoody manifolds, with cyclically presented fundamental groups. It is proved in [15] and [7] that the family of Dunwoody manifolds coincides with the family of strongly-cyclic coverings of lens spaces (possibly $\mathbf{S}^3$), branched over $(1, 1)$ -knots. As a consequence, any $(1, 1)$ -knot can be represented by four integers a, b, c, r , and it will be denoted by $K(a, b, c, r)$.

In this paper we show that the orientable Seifert manifold with invariants

$$\{Oo, 0 \mid -1; \underbrace{(p, q), \dots, (p, q)}_{n \text{ times}}, (l, l-1)\}$$

has the fundamental group isomorphic to the cyclically presented group $G_n((x_1^q \cdots x_n^q)^l x_n^{-p})$, and it is the n -fold strongly-cyclic covering of the lens space $L(|nlq-p|, q)$, branched over the $(1, 1)$ -knot $K(q, q(nl-2), p-2q, p-q)$, if $p \geq 2q$, and over the $(1, 1)$ -knot $K(p-q, 2q-p, q(nl-2), p-q)$, if $p < 2q$.

2 Basic notions

A finite balanced presentation of a group $\langle x_1, \dots, x_n \mid r_1, \dots, r_n \rangle$ is said to be a *cyclic presentation* if there exists a word w in the free group F_n generated by $x_1, \dots, x_n$ such that $r_k = \theta^{k-1}(w)$, $k = 1, \dots, n$, where $\theta : F_n \rightarrow F_n$ denotes the automorphism defined by $\theta(x_i) = x_{i+1}$ (subscripts mod n), $i = 1, \dots, n$. This presentation (and the related group) will be denoted by $G_n(w)$. For further details see [18].

A knot K in a closed, connected, orientable 3-manifold N^3 is called a $(1, 1)$ -knot if there exists a Heegaard splitting of genus one $(N^3, K) = (H, A) \cup_\varphi (H', A')$, where H and H' are solid tori, $A \subset H$ and $A' \subset H'$ are properly embedded trivial arcs¹, and $\varphi : (\partial H', \partial A') \rightarrow (\partial H, \partial A)$ is an attaching homeomorphism (see Figure 1). Obviously, N^3 turns out to be a lens space $L(p, q)$, including $\mathbf{S}^3 = L(1, 0)$ and $\mathbf{S}^2 \times \mathbf{S}^1 = L(0, 1)$.

¹This means that there exists a disk $D \subset H$ (resp. $D' \subset H'$) with $A \cap D = A \cap \partial D = A$ and $\partial D - A \subset \partial H$ (resp. $A' \cap D' = A' \cap \partial D' = A'$ and $\partial D' - A' \subset \partial H'$).




Figure 1: A $(1,1)$ -knot decomposition.

It is well known that the family of $(1,1)$ -knots contains all torus knots and all two-bridge knots in $\mathbf{S}^3$. Several topological properties of $(1,1)$ -knots have recently been investigated in many papers (see references in [6]).

An integer 4-parametric representation of $(1,1)$ -knots have been developed in [7] (see also [8]). Every $(1,1)$ -knot, with the only exception of the “core” knot $\{P\} \times \mathbf{S}^1 \subset \mathbf{S}^2 \times \mathbf{S}^1$, can be represented by four non-negative integers a, b, c, r , and the represented knot will be referred as $K(a, b, c, r)$.

A $(1,1)$ -knot $K(a, b, c, r)$, with $a + b + c > 0$, admits a natural $(1,1)$ -decomposition $(H, A) \cup_{\varphi} (H', A')$ described by the genus one Heegaard diagram of Figure 2, where the labels a, b and c denote the correspondent number of parallel arcs, and the gluing between the circles C' and C'' depends on the twist parameter r in such a way that equally labelled vertices are identified together (observe that r can be taken mod $2a + b + c$).

As a simple consequence of Seifert-Van Kampen Theorem, the fundamental group of the exterior of a $(1,1)$ -knot (as well as its first homology group) is generated by the two loops $\alpha, \gamma \subset \partial H$ depicted in Figure 3.

In the next section we need the following result.

Lemma 1 (i) *If a and c are non-negative integers such that $\gcd(a, c) = 1$, then $K(a, 0, c, a)$ is a $(1,1)$ -knot in the lens space $L(c, a)$.*

(ii) *If a, b, c are non-negative integers such that $\gcd(a - c, b + c) = 1$, then $K(a, b, c, a)$ is a $(1,1)$ -knot in the lens space $L(b + c, a + b)$.*

(iii) *If a, b, c are non-negative integers such that $a > 0$ and $\gcd(a, b - c) = 1$, then $K(a, b, c, a + c)$ is a $(1,1)$ -knot in the lens space $L(|b - c|, a)$.*

Proof. As proved in [8], $K(a, b, c, r)$ is equivalent to $K(a, c, b, 2a + b + c - r)$ and $K(a, 0, c, r)$ is equivalent to $K(a, c, 0, r)$. As a consequence, $K(a, 0, c, r)$ is equivalent to $K(a, 0, c, 2a + c - r)$.




Figure 2: Heegaard diagram for $K(a, b, c, r)$




Figure 3:

(i) If $a = 0$, then $c = 1$ and the result is straightforward. If $0 < a < c$, then applying the Singer move [29] of type IB depicted in Figure 4, we obtain the canonical Heegaard diagram of $L(c, a)$. If $a \geq c$, the Singer move depicted in Figure 5 transforms the diagram of $K(a, 0, c, a)$ in the diagram of $K(a - c, c, 0, a - c)$, which is equivalent to $K(a - c, 0, c, a - c)$. Since $L(c, a - c)$ is homeomorphic to $L(c, a)$, the result follows from the previous case $a < c$.

(ii) If $b = 0$ the result follows from (i). If $b > 0$, by performing the Singer move of Figure 6, the diagram of $K(a, b, c, a)$ becomes the diagram of $K(a - 1, b + 1, c - 1, a - 1)$. If $a \leq c$, after performing the move a times, we obtain the diagram of $K(0, b + a, c - a, 0)$, which is the canonical Heegaard diagram of $L(b + c, a + b)$, since $\gcd(a + b, b + c) = \gcd(a - c, b + c) = 1$. If $a > c$, after performing the move c times, we obtain the diagram of $K(a - c, b + c, 0, a - c)$, which is equivalent to $K(a - c, 0, b + c, a - c)$. Now the result follows from (i), since $L(b + c, a - c)$ is homeomorphic to $L(b + c, a + b)$.

(iii) Since $K(a, b, c, a + c)$ is equivalent to $K(a, c, b, a + b)$, we can always suppose $c \leq b$. If $c > 0$, then, by performing the Singer move of Figure 7, the diagram of $K(a, b, c, a + c)$ becomes the diagram of $K(a, b - 1, c - 1, a + c - 1)$. After performing the move c times, we obtain the diagram of $K(a, b - c, 0, a)$, which is equivalent to $K(a, 0, b - c, a)$. The result now follows from (i). ■

Observe that, since $K(a, b, c, a + b + c)$ is equivalent to $K(a, c, b, a)$, $K(a, b, c, a + b + c)$ is a $(1, 1)$ -knot in the lens space $L(b + c, a + c)$, if $\gcd(a - b, b + c) = 1$.

An n -fold cyclic covering M^3 of a 3-manifold N^3 , branched over a knot $K \subset N^3$, is called *strongly-cyclic* if the branching index of K is n . This means that the fiber in M^3 of each point of K consists of a single point. In this case, the homology class m of a meridian loop around K is mapped by the monodromy $\omega : H_1(N^3 - K) \rightarrow \mathbf{Z}_n$ of the covering in a generator of $\mathbf{Z}_n$ (up to equivalence, we can always suppose $\omega(m) = 1$). Observe that a cyclic branched covering of a knot K in $\mathbf{S}^3$ is always strongly-cyclic and is uniquely determined, up to equivalence, since $H_1(\mathbf{S}^3 - K) \cong \mathbf{Z}$. Obviously, this property is no longer true for a knot in a more general 3-manifold. Necessary and sufficient conditions for the existence and uniqueness of strongly-cyclic branched coverings of $(1, 1)$ -knots have been obtained in [5].




Figure 4:




Figure 5:




Figure 6:




Figure 7:

3 Main results

Let n, p, q, l be positive integers such that $q < p$ and $\gcd(p, q) = 1$. In the following we denote by $\Sigma(n, p, q, l)$ the orientable Seifert manifold [28] with invariants

$$\{Oo, 0 \mid -1; \underbrace{(p, q), \dots, (p, q)}_{n \text{ times}}, (l, l-1)\}$$

having $\mathbf{S}^2$ as orbit space, n exceptional fibres of type (p, q) and, for $l > 1$, an exceptional fibre of type $(l, l-1)$.

Observe that, in particular, the manifolds $\Sigma(n, 2, 1, 1)$ are precisely the Neuwirth manifolds M_n introduced in [25], studied in [9], and generalized in [16], [31] and [27].

Proposition 2 *The fundamental group of $\Sigma(n, p, q, l)$ is cyclically presented by $G_n(w)$, where $w = (x_1^q \cdots x_n^q)^l x_n^{-p}$.*

Proof. Following [26], a standard presentation of $G = \pi_1(\Sigma(n, p, q, l))$ is

$$\langle y_1, \dots, y_n, y, h \mid [y_i, h], [y, h], y_i^p h^q, y^l h^{l-1}, y_1 \cdots y_n y h; i = 1, \dots, n \rangle.$$

Since $\gcd(p, q) = 1$, there exist $\alpha, \beta \in \mathbf{Z}$ such that $q\beta - p\alpha = 1$.

From the relations we have $y^l h^{l-1} = (yh)^l h^{-1}$. By introducing the new generator $x = yh$, the presentation becomes

$$\langle y_1, \dots, y_n, x, h \mid [y_i, h], [x, h], y_i^p h^q, x^l h^{-1}, y_1 \cdots y_n x; i = 1, \dots, n \rangle.$$

Now we define $x_i = y_i^\beta h^\alpha$, for $i = 1, \dots, n$. Then we have the following new presentation for G :

$$\langle y_1, \dots, y_n, x, h, x_1, \dots, x_n \mid [y_i, h], [x, h], y_i^p h^q, x^l h^{-1}, y_1 \cdots y_n x, x_i^{-1} y_i^\beta h^\alpha; i = 1, \dots, n \rangle.$$

From the relations we obtain $x_i^q = y_i^{q\beta} h^{q\alpha} = y_i^{1+p\alpha} h^{q\alpha} = y_i (y_i^p h^q)^\alpha = y_i$ and $x_i^p = y_i^{p\beta} h^{p\alpha} = y_i^{p\beta} h^{q\beta-1} = (y_i^p h^q)^\beta h^{-1} = h^{-1}$.

Therefore G admits the presentation

$$\langle x, h, x_1, \dots, x_n \mid [x_i^q, x_i^{-p}], [x, x^l], x_i^{qp} x_i^{-qp}, x^l h^{-1}, x_1^q \cdots x_n^q x, x_i^{-1} x_i^{q\beta} x_i^{-p\alpha}, x_i^p h; i = 1, \dots, n \rangle,$$

which obviously becomes

$$\langle x, h, x_1, \dots, x_n \mid x^l h^{-1}, x_1^q \cdots x_n^q x, x_i^p h; i = 1, \dots, n \rangle.$$

From the relations we obtain $x = (x_1^q \cdots x_n^q)^{-1}$ and $h = (x_1^q \cdots x_n^q)^{-l}$.

Therefore G admits the presentation:

$$\langle x_1, \dots, x_n \mid (x_1^q \cdots x_n^q)^l x_i^{-p}; i = 1, \dots, n \rangle,$$

which is equivalent to

$$\langle x_1, \dots, x_n \mid (x_1^q \cdots x_n^q)^l x_n^{-p}, x_i^p x_{i+1}^{-p}; i = 1, \dots, n-1 \rangle.$$

In order to prove that G is isomorphic to $G_n(w)$, it suffices to show that the normal closures in the free group F_n of the sets $\{w, x_i^p x_{i+1}^{-p}; i = 1, \dots, n-1\}$ and $\{w, \theta^i(w); i = 1, \dots, n-1\}$ coincide.

This is achieved by observing that, setting $w_i = \theta^i(w)$, we have:

$$w_i^{-1} x_{i+1}^q w_{i+1} x_{i+1}^{-q} = x_i^p x_{i+1}^{-p}, \text{ for } i = 1, \dots, n-2,$$

$$w_{n-1}^{-1} x_n^q w x_n^{-q} = x_{n-1}^p x_n^{-p},$$

$$x_1^{-q} w ((x_1^p x_2^{-p}) \cdots (x_{n-1}^p x_n^{-p}))^{-1} x_1^q = w_1,$$

$$x_i^{-q} w_{i-1} (x_{i-1}^p x_i^{-p}) x_i^q = w_i, \text{ for } i = 2, \dots, n-1.$$

■

Since a Seifert manifold with one or two singular fibers is trivially a lens space (see [26]), we will suppose $n > 1$ and $l > 1$ when $n = 2$.

Theorem 3 *The Seifert manifold $\Sigma(n, p, q, l)$ is the n -fold strongly-cyclic covering of the lens space $L(|nlq - p|, q)$, branched over the $(1, 1)$ -knot $K(q, q(nl - 2), p - 2q, p - q)$, if $p \geq 2q$, and over the $(1, 1)$ -knot $K(p - q, 2q - p, q(nl - 2), p - q)$, if $p < 2q$.*

Proof. (i) Suppose $p \geq 2q$. By (iii) of Lemma 1, $K = K(q, q(nl - 2), p - 2q, p - q)$ is a $(1, 1)$ -knot in $L(|nlq - p|, q)$. Now suppose that K admits the n -fold strongly-cyclic branched covering $C_n(K)$ defined by $\omega(\alpha) = 0$ and $\omega(\gamma) = 1$. In this case $C_n(K)$ is the Dunwoody manifold $D(q, q(nl - 2), p - 2q, n, p - q, 0)$ (see [15] and [7]). Its defining genus n Heegaard diagram, having cyclic symmetry of order n , is depicted in Figure 8, where $a = q, b = q(nl - 2), c = p - 2q, r = p - q$, and the circle C'_i must be glued to the circle C''_i , for $i = 1, \dots, n$, according to the twist parameter r . In order to check if the above diagram really represents a manifold, it is convenient to consider the cellular decomposition dual to the one associated to the diagram. In this way, $C_n(K)$ is obtained by pairwise identification of the regions of the 2-cell tessellation of the boundary of a 3-ball, as depicted in Figure 9. The tessellation consists of $2n$ regions, $R'_1, \dots, R'_n$ around the North pole N and

$R_1'', \dots, R_n''$ around the South pole S , corresponding to the cycles C_i' and C_i'' of the diagram. The edges of the tessellation correspond to the arcs of the diagram in the following way. The poles S and N are connected by n meridians and each meridian m_i is composed by nlq edges $e_{i,j}$ ($j = 1, \dots, nlq$) from S to N , and precisely q from S to B_i , $q(nl - 2)$ from B_i to A_i and q from A_i to N . Moreover, for $i = 1, \dots, n$, the arc a_i connecting B_{i-1} with A_i is composed by $p - 2q$ edges $e'_{i,j}$ ($j = 1, \dots, p - 2q$). Observe that, when $p = 2q$ (i.e., $p = 2$ and $q = 1$), the points B_{i-1} and A_i coincide. In order to obtain $C_n(K)$, the region R_i' is glued to R_i'' , for $i = 1, \dots, n$, by an orientation reversing homeomorphism in such a way that point S of R_i'' matches point² B_{i-1} of R_i' . By this gluing we have: $e_{i,j} \equiv e_{i-1,j+q}$ for $j = 1, \dots, q(nl - 1)$, $e_{i-1,j} \equiv e'_{i,j}$ for $j = 1, \dots, q$, $e'_{i,j} \equiv e'_{i,j+q}$ for $j = 1, \dots, p - 3q$, and $e'_{i,p-3q+k} \equiv e_{i,q(nl-1)+k}$ for $k = 1, \dots, q$. As a consequence of all identifications, we have $e_{i,j} \equiv e_{i,j+nq} \equiv \dots \equiv e_{i,j+n(l-1)q}$ for $j = 1, \dots, nq$, and moreover $e_{i-1,j} \equiv e_{i,q(nl-1)+j}$ for $j = 1, \dots, q$. Since $\gcd(p, q) = 1$, all edges of the arcs SB_{i-1} , $B_{i-1}A_i$ and A_iN match, and we will call them x_i . Therefore, the $q(nl - 2)$ edges of the arc B_iA_i become: q times x_{i+1} , q times x_{i+2} , $\dots$, q times x_i ($l - 1$ times) and then q times x_{i+1} , q times x_{i+2} , $\dots$, q times x_{i-2} (see Figure 10). Since each edge x_i appears p times consecutively, its endpoints coincide. Moreover, since every edge x_i has S as endpoint, the cellular decomposition of $C_n(K)$ is composed by one vertex, n edges, n regions and one 3-ball. By Seifert criterion, $C_n(K)$ is actually a closed, orientable 3-manifold.

Moreover, $\pi_1(C_n(K))$ admits a balanced presentation with $x_1, \dots, x_n$ as generators and with relators obtained by walking around the boundaries of the regions R_i' . As a consequence, $\pi_1(C_n(K)) = G_n((x_1^q \dots x_n^q)^l x_n^{-p})$, and therefore it is isomorphic to $\pi_1(\Sigma(n, p, q, l))$.

In order to prove that $C_n(K)$ and $\Sigma(n, p, q, l)$ are actually homeomorphic, we note that $\pi_1(C_n(K))$ has non-trivial centre. So, either $C_n(K)$ is prime or is the connected sum $C_n(K) = M \# M'$, where $\pi_1(M')$ is trivial and M is prime, with $\pi_1(M) = \pi_1(C_n(K))$.

If $\Sigma(n, p, q, l)$ is large (in the sense of [26]), then also M is a large Seifert manifold (see [4]). So M and $\Sigma(n, p, q, l)$ are homeomorphic, with Heegaard genus at least $n - 2$ (see [2]). If M' is not homeomorphic to $\mathbf{S}^3$, then its Heegaard genus is greater than two, and $C_n(K)$ has Heegaard genus greater than n . But this is impossible since $C_n(K)$ admits a genus n Heegaard

²The subscript i is considered mod n .

splitting by [24]. As a consequence, $M' = \mathbf{S}^3$ and $C_n(K) = \Sigma(n, p, q, l)$.

If $\Sigma(n, p, q, l)$ is not large, then one of the following possibilities holds: (a) $n = 3, p = 2, q = 1 = l$; (b) $n = 2, p = 2, q = 1, l > 1$; (c) $n = 2, p = 3, q = 1, l = 2$. Since for $q = 1$ the given decomposition of $C_n(K)$ coincides with the decomposition $P(p, \dots, p; l)$ given in [27], the result follows from [27, Proposition 4.1].

(ii) Let $p < 2q$. By (ii) of Lemma 1, $K(p - q, 2q - p, q(nl - 2), p - q)$ is a $(1, 1)$ -knot in $L(|nlq - p|, q)$. Now suppose that K admits the n -fold strongly-cyclic branched covering $C_n(K)$ defined by $\omega(\alpha) = 1$ and $\omega(\gamma) = 1$. In this case $C_n(K)$ is the Dunwoody manifold $D(p - q, 2q - p, q(nl - 2), n, p - q, 1)$. Its defining genus n Heegaard diagram, having cyclic symmetry of order n , is depicted in Figure 8, where $a = p - q, b = 2q - p, c = q(nl - 2), r = p - q$, and the circle C'_i must be glued to the circle C''_{i+1} , for $i = 1, \dots, n$, according to the twist parameter r . Again it is better to refer to the dual decomposition (see Figure 9). This time each meridian is composed by p edges, and precisely $p - q$ from S to B_i , $2q - p$ from B_i to A_i and $p - q$ from A_i to N . Moreover, the arc connecting B_{i-1} with A_i is composed by $q(nl - 2)$ edges. In order to obtain $C_n(K)$, the region R'_{i-1} is glued to R''_i in such a way that point N of R'_{i-1} matches point B_{i-1} of R''_i .

In complete analogy with point (i), it is easy to see that the gluing gives rise to a cellular decomposition of $C_n(K)$, composed by one vertex, n edges, n regions and one 3-ball (see the 2-skeleton in Figure 11). So $\pi_1(\Sigma(n, p, q, l))$ is isomorphic to $\pi_1(C_n(K))$ and $\Sigma(n, p, q, l)$ is homeomorphic to $C_n(K)$ when it is large. The only small Seifert manifold occurs when $n = q = l = 2, p = 3$. To achieve the result, it suffices to prove that $D = D(1, 1, 4, 2, 1, 1)$ is the 2-fold covering of $\mathbf{S}^3$, branched over the Montesinos knot $\mathbf{m}(-1; 1/2, 2/3, 2/3)$ (see [3, Chapter 12]). A genus two Heegaard diagram of D is depicted in Figure 12, where the circle C'_1 (resp. C'_2) must be glued to the circle C''_2 (resp. C''_1) in such a way that equally labelled vertices match. The application of the Takahashi algorithm [32] to this diagram (as depicted in Figure 13) shows that the manifold D is the 2-fold branched covering of the knot K represented by the 3-bridge diagram of Figure 14. A Wirtinger presentation of the fundamental group of the exterior of K can be easily computed from this diagram, obtaining $\pi_1(\mathbf{S}^3 - K) = \langle x, y, z \mid yz^{-1}xzy^{-1}xyzy^{-1}x^{-1}zy^{-1}z^{-1}, x^{-1}zyzy^{-1}xzx^{-1}zy^{-1}z^{-1}xy^{-1} \rangle$. The Alexander polynomial of K can be obtained by a standard application of Fox differential calculus (see [3, Chapter 9]), and we have $\Delta_K(t) = 1 - 4t + 5t^2 - 4t^3 + t^4$. By the sequence of Reidemeister moves depicted in Figures 15-17, the knot K





Figure 8:



admits the 9 crossing diagram of Figure 17. Since the knot 8_{21} of Rolfsen's table is the only knot with crossing number ≤ 9 with Alexander polynomial $1 - 4t + 5t^2 - 4t^3 + t^4$ (see [19, Table 3]), then K is precisely the knot 8_{21} . Since 8_{21} is the Montesinos knot $\mathbf{m}(-1; 1/2, 2/3, 2/3)$ (see [19, Table 2], which uses a slightly different notation), the statement is proved. ■

Observe that Theorem 3 refines and extends the results of [13, Theorem 4.2] and [30, Theorem 5.1]. The proof of Theorem 3 also gives the following:

Corollary 4 *The Seifert manifold $\Sigma(n, p, q, l)$ is the Dunwoody manifold $D(q, q(nl - 2), p - 2q, n, p - q, 0)$ if $p \geq 2q$ and the Dunwoody manifold $D(p - q, 2q - p, q(nl - 2), n, p - q, 1)$ if $p < 2q$.*

Remark Since $\Sigma(n, p, p - 1, 1) = \Sigma(n - 1, p, p - 1, p)$, this manifold is at the same time the n -fold strongly-cyclic covering of the lens space $L(pn - p - n, p - 1)$, branched over the $(1, 1)$ -knot $K(1, p - 2, (p - 1)(n - 2), 1)$ and the $(n - 1)$ -fold strongly-cyclic covering of the lens space $L(p(pn - p - n), p - 1)$, branched over the $(1, 1)$ -knot $K(1, p - 2, (p - 1)(np - p - 2), 1)$, when $p > 2$. Moreover, the Neuwirth manifold $M_n = \Sigma(n, 2, 1, 1) = \Sigma(n - 1, 2, 1, 2)$ is at the same time the n -fold strongly-cyclic covering of the lens space $L(n - 2, 1)$, branched over the $(1, 1)$ -knot $K(1, n - 2, 0, 1)$ and the $(n - 1)$ -fold strongly-cyclic covering of the lens space $L(2n - 4, 1)$, branched over the $(1, 1)$ -knot $K(1, 2n - 4, 0, 1)$.








Figure 13:




Figure 14:




Figure 15:




Figure 16:




Figure 17:

Acknowledgements

Work performed under the auspices of the G.N.S.A.G.A. of I.N.d.A.M. (Italy) and the University of Bologna, funds for selected research topics.

References

- [1] P. Bandieri, A. C. Kim and M. Mulazzani: *On the cyclic coverings of the knot 5_2* . Proc. Edinb. Math. Soc. **42** (1999), 575–587.
- [2] M. Boileau and H. Zieschang: *Heegaard genus of closed orientable Seifert 3-manifolds*. Invent. Math. **76** (1984), 455–468.
- [3] G. Burde and H. Zieschang: *Knots*. De Gruyter Studies in Mathematics, 5. Walter de Gruyter, 1985.
- [4] A. Casson and D. Jungreis: *Convergence groups and Seifert fibered 3-manifolds*. Invent. Math. **118** (1994), 441–456.
- [5] A. Cattabriga and M. Mulazzani: *Strongly-cyclic branched coverings of $(1, 1)$ -knots and cyclic presentation of groups*. Math. Proc. Camb. Philos. Soc. **135** (2003), 137–146.

- [6] A. Cattabriga and M. Mulazzani: *(1, 1)-knots via the mapping class group of the twice punctured torus*. Adv. Geom. **4** (2004), 263–277.
- [7] A. Cattabriga and M. Mulazzani: *All strongly-cyclic branched coverings of $(1, 1)$ -knots are Dunwoody manifolds*. J. London Math. Soc. **70** (2004), 512–528.
- [8] A. Cattabriga and M. Mulazzani: *Representations of $(1, 1)$ -knots*. to appear in Fundamenta Mathematicae, Proceedings of the Second International Conference “Knots in Poland 2003”, Vol. II (2005). arXiv:math.GT/0501234
- [9] A. Cavicchioli: *Neuwirth manifolds and colourings of graphs*. Aequationes Math. **44** (1992), 168–187.
- [10] A. Cavicchioli, F. Hegenbarth and A. C. Kim: *A geometric study of Sieradsky groups*. Algebra Colloq. **5** (1998), 203–217.
- [11] A. Cavicchioli, F. Hegenbarth and A. C. Kim: *On cyclic branched coverings of torus knots*. J. Geom. **64** (1999), 55–66.
- [12] A. Cavicchioli, F. Hegenbarth and D. Repovs: *On manifold spines and cyclic presentations of groups*. In: Knot theory. Proceedings of the mini-semester, Warsaw, Poland, July 13–August 17, 1995. Warszawa: Polish Academy of Sciences, Institute of Mathematics, Banach Cent. Publ. **42** (1998), 49–56.
- [13] A. Cavicchioli, D. Repovs and F. Spaggiari: *Topological properties of cyclically presented groups*. J. Knot Theory Ramifications **12** (2003), 243–268.
- [14] M. J. Dunwoody: *Cyclic presentations and 3-manifolds*. In: Proc. Inter. Conf., Groups-Korea '94, Walter de Gruyter, Berlin-New York (1995), 47–55.
- [15] L. Grasselli and M. Mulazzani: *Genus one 1-bridge knots and Dunwoody manifolds*. Forum Math. **13** (2001), 379–397.
- [16] L. Grasselli and S. Piccarreta: *Crystallizations of generalized Neuwirth manifolds*. Forum Math. **9** (1997), 669–685.

- [17] H. Helling, A. C. Kim and J. L. Mennicke: *A geometric study of Fibonacci groups*. J. Lie Theory **8** (1998), 1-23.
- [18] D. L. Johnson: *Topics in the theory of group presentations*. London Math. Soc. Lect. Note Ser., vol. 42, Cambridge Univ. Press, Cambridge, U.K., 1980.
- [19] A. Kawauchi: *A survey of knot theory*. Birkhauser Verlag, Basel, 1996.
- [20] A. C. Kim: *On the Fibonacci group and related topics*. Contemp. Math. **184** (1995), 231–235.
- [21] A. C. Kim, Y. Kim and A. Vesnin: *On a class of cyclically presented groups*. In: Proc. Inter. Conf., Groups-Korea '98, Walter de Gruyter, Berlin-New York (2000), 211–220.
- [22] G. Kim, Y. Kim and A. Vesnin: *The knot 5_2 and cyclically presented groups*. J. Korean Math. Soc. **35** (1998), 961–980.
- [23] C. Maclachlan and A. W. Reid: *Generalised Fibonacci manifolds*. Transform. Groups **2** (1997), 165–182.
- [24] M. Mulazzani: *Cyclic presentations of groups and cyclic branched coverings of $(1, 1)$ -knots*. Bull. Korean Math. Soc. **40** (2003), 101–108.
- [25] L. Neuwirth: *An algorithm for the construction of 3-manifolds from 2-complexes*. Proc. Camb. Philos. Soc. **64** (1968), 603–613.
- [26] P. Orlik: *Seifert manifolds*. Lecture Notes in Mathematics, Vol. 291. Springer-Verlag, Berlin-New York, 1972.
- [27] B. Ruini, F. Spaggiari and A. Vesnin: *On spines of Seifert fibered manifolds*. Aequationes Math. **65** (2003), 40–60.
- [28] H. Seifert and W. Threlfall: *Seifert and Threlfall: a textbook of topology*. Pure and Applied Mathematics, 89. Academic Press, New York-London, 1980.
- [29] J. Singer: *Three-dimensional manifolds and their Heegaard diagrams*. Trans. Amer. Math. Soc. **35** (1933), 88-111.

- [30] F. Spaggiari: *The combinatorics of some Tetrahedron manifolds*. Discrete Math., to appear.
- [31] A. Szczepański and A. Vesnin: *Generalized Neuwirth groups and Seifert fibered manifolds*. Algebra Colloq. **7** (2000), 295–303.
- [32] M. Takahashi: *Two knots with the same 2-fold branched covering space*. Yokohama Math. J. **25** (1977), 91-99.
- [33] A. Vesnin and A. C. Kim: *The fractional Fibonacci groups and manifolds*. Sib. Math. J. **39** (1998), 655–664.

LUIGI GRASSELLI, Department of Sciences and Methods for Engineering, University of Modena and Reggio Emilia, 42100 Reggio Emilia, ITALY. E-mail: grasselli.luigi@unimore.it

MICHELE MULAZZANI, Department of Mathematics, University of Bologna, I-40127 Bologna, ITALY, and C.I.R.A.M., Bologna, ITALY. E-mail: mulazza@dm.unibo.it